



Joshua Henry

Brian E. Whipker
bwhipker@ncsu.edu

Revising Your Phosphorus Fertilization Strategy

by Joshua Henry and Brian E. Whipker

Results of current research projects being conducted at NC State University are reported here.

Phosphorus fertilization strategies for greenhouse crops are in flux. Fertilizer formulations that have been used for many years actually supply far more phosphorus than most crops need. In fact, only 5 to 10 ppm is usually enough for healthy, compact growth in most greenhouse crops. Given this range, it is apparent that fertilizers such as 20-20-20 and 20-10-20 supply 10 to 15 times more phosphorus than necessary even at recommended rates (Figure 1).

Research is currently being conducted at North Carolina State University on phosphorus. In one experiment, we are determining the plant growth response to various levels of phosphorus fertilization. Trials have been done on *Alternanthera* and *Iresine* species to determine optimal phosphorus levels. These trials were conducted using rates between 0 and 80 ppm to create a visual growth response curve (Figures 2&3).

Figure 1. Phosphorus Content of Common Fertilizers.

Fertilizer Formulation	Phosphorus Levels (ppm)	
	Mixed at 100 ppm N	Mixed at 200 ppm N
20-20-20	43	83
20-10-20	21.5	43
15-5-15 Cal Mag	10.75	21.5
13-2-13 Cal Mag	4.3	8.6
15-0-15	0	0

Summary of Findings

- Most fertilization programs provide phosphorus in excess of the plant's needs.
- For the majority of plants grown in the greenhouse, phosphorus levels of 5 to 10 ppm are adequate.
- Zero phosphorus fertilization practices should be avoided due to the onset of deficiency symptoms and the potential for crop losses.
- Phosphorus deficiency symptoms develop in two ways. The most common symptom is lower leaf purpling. The second type of symptom begins as an olive-green spotting on the lower leaves and occurs during warmer growing conditions.
- Keep in mind that root rot, excessive irrigations, and cold growing can also limit phosphorus uptake. So it is always a good idea to inspect the root system.

When no phosphorus is supplied to the plant, it will have stunted growth and remain quite compact. As small increments of phosphorus are added to the fertilizer solution, growth rapidly increases up to about 5 to 10 ppm depending on the species. After this point, growth begins to plateau and remains fairly constant as more phosphorus is added. This plateau goes on for some time, but growth will eventually begin to decrease due to toxic levels of phosphorus building up in the plant tissue. This decrease in growth affects both height and dry weight, and can begin to occur at rates above 40 to 80 ppm phosphorus (Figure 4).

Additional research has been conducted by Dr. Paul Nelson and his graduate students at NCSU over the past several decades. In his research, it has been found that reduced or even 0 ppm phosphorus regiments can be very beneficial for plug production by limiting stretch (Figure 5). Of course, no phosphorus can only be used for short term crops such as plugs, as deficiency symptoms can quickly develop when plants do not have adequate phosphorus. In 2014, there were two major cases where losses occurred with garden mum crops when phosphorus was limited (see e-GRO 3-62).

Deficiency symptoms related to phosphorus often include the typical reddening or purpling of the lower foliage (Figure 6). Olive green spots can occur as another deficiency symptom as well, especially during warmer growing conditions. These spots often develop within a chlorotic area in the center of the leaf (Figure 7).

It is important to note that phosphorus deficiency can occur even when sufficient rates are supplied in the fertilizer. Certain growing conditions can greatly affect the plants ability to take up phosphorus and can result in deficiency symptoms. Issues with root health including root rot are one set of conditions which can lead to phosphorus deficiency

Figure 2. Phosphorus Response Curve for Iresine.






Figure 3. Phosphorus Response Curve for Alternanthera.



(Figure 8). Cold and wet conditions as well as drought stress can also lead to a deficiency situation (Figure 9). In all of these cases, plants lack the ability to take up adequate phosphorus and can therefore exhibit the red or purple discoloration often associated with phosphorus deficiency.

Reddening of the lower leaves can however occur for a number of unrelated reasons. These “mimics” can be due to low substrate pH, low EC or even sulfur deficiency. Due to

Figure 4. *Iresine* Dry Weight and Height Response to Increasing P Rates.



Treatment Key (in ppm P): 0, 1.25, 2.5, 5, 10, 20 40, Or 80.
Henry, unpublished data



Reducing phosphorus fertilization rates can have a number of benefits for growers such as reducing fertilizer costs. Phosphorus runoff has also been an issue affecting water quality and causing algae blooms such as those in Lake Erie. Reduction of runoff has definite benefits for the environment. Additionally, using a low phosphorus fertilization strategy can produce more compact plants. When plants are more compact, growers can benefit from a potential reduction in spacing during production, as well as fitting more shelves per shipping rack. Lastly, compact plants are generally more desirable from the consumer standpoint. There are numerous reasons to revise your phosphorus fertilization strategy, so this hopefully this information will help in doing so.

the variety of potential causes, conducting pour-through tests or even tissue analysis are important first steps to take to determine the issue.

Symptoms of toxicity are less likely to occur, but it is still good to recognize them if high levels of phosphorus are being supplied to the crop. Toxicity symptoms manifest themselves as iron deficiency, or interveinal chlorosis of the upper foliage. When phosphorus reaches toxic levels, it has an antagonistic effect on iron uptake by the plant. This is most common in crops native to Australia such as scaevola, as they are highly sensitive to elevated phosphorus levels. Deficiency and toxicity alike can usually be avoided by providing the optimal range of 5 to 10 ppm phosphorus.

Low phosphorus fertilization regimens can be highly beneficial for producing healthy ornamental crops in the greenhouse setting. The optimal range of phosphorus is about 5 to 10 ppm. These rates can be achieved using 13-2-13 Cal Mag as a constant feed mixed at 100 or 200 ppm nitrogen. Using 15-5-15 Cal Mag as a constant feed at 100 ppm nitrogen will also supply sufficient levels. A third option could involve alternating between 15-5-15 at 200 ppm nitrogen and 15-0-15 to achieve the same effect.

Figure 5. Utilizing a Low Phosphorus Strategy on Plugs for Compactness.

P₂O₅ as a % of N in post-plant fertilizer



Source: Paul Nelson, NCSU

P₂O₅ = 50% of N
(e.g. 20-10-20)

15%
(13-2-13)

7.5%

0%



Figure 6. Phosphorus Deficiency:
Lower Leaf - Red Coloration



Whipker, unpublished data

Figure 7. Phosphorus Deficiency:
Lower Leaf - Olive Green Leaf Spots



Whipker, unpublished data

Figure 8. Phosphorus Deficiency:
Lower Leaf - Red Coloration



Limited P uptake due to root rot



Whipker, unpublished data

Figure 9. Phosphorus Deficiency:
Lower Leaf - Red Coloration



Limited P uptake due to cold/wet conditions



Whipker, unpublished data

© Brian Whipker, 2015. No reproduction please.